

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07C 405/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 98/21182</b> <b>(43) International Publication Date:</b> 22 May 1998 (22.05.98)
<b>(21) International Application Number:</b> PCT/US97/20857 <b>(22) International Filing Date:</b> 7 November 1997 (07.11.97)  <b>(30) Priority Data:</b> 60/030,504 12 November 1996 (12.11.96) US  <b>(71) Applicant (for all designated States except US):</b> ALCON LABORATORIES, INC. [US/US]; 6201 South Freeway, Fort Worth, TX 76134-2099 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> KLIMKO, Peter, G. [US/US]; 2115 Pembroke Drive, Fort Worth, TX 76110 (US). ZINKE, Paul, W. [US/US]; 4129 Willow Way Road, Fort Worth, TX 76133 (US).  <b>(74) Agents:</b> COPELAND, Barry, L. et al.; Alcon Laboratories, Inc., Patent Dept., Q-148, 6201 South Freeway, Fort Worth, TX 76134-2099 (US).		<b>(81) Designated States:</b> AU, CA, CN, JP, KR, MX, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> <i>CIS-<math>\Delta^4</math> ANALOGS OF PROSTAGLANDINS AS OCULAR HYPOTENSIVES</i>  <b>(57) Abstract</b>  <i>Cis-<math>\Delta^4</math> analogs of prostaglandins and methods of their use in treating glaucoma and ocular hypertension are disclosed.</i>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

***CIS- $\Delta^4$*  ANALOGS OF PROSTAGLANDINS  
AS OCULAR HYPOTENSIVES**

**Background of the Invention**

The present invention relates to novel compounds for the treatment of glaucoma and ocular hypertension. In particular, the present invention relates to the use of certain *cis- $\Delta^4$*  analogs of D and F series prostaglandins to treat glaucoma and ocular hypertension.


Glaucoma is a progressive disease which leads to optic nerve damage, and, ultimately, total loss of vision. The causes of this disease have been the subject of extensive studies for many years, but are still not fully understood. The principal symptom of and/or risk factor for the disease is elevated intraocular pressure or ocular hypertension due to excess aqueous humor in the anterior chamber of the eye.

The causes of aqueous humor accumulation in the anterior chamber are not fully understood. It is known that elevated intraocular pressure ("IOP") can be at least partially controlled by administering drugs which either reduce the production of aqueous humor within the eye, such as beta-blockers and carbonic anhydrase inhibitors, or increase the flow of aqueous humor out of the eye, such as miotics and sympathomimetics.

Most types of drugs conventionally used to treat glaucoma have potentially serious side effects. Miotics such as pilocarpine can cause blurring of vision and other visual side effects, which may lead either to decreased patient compliance or to termination of therapy. Systemically administered carbonic anhydrase inhibitors can also cause serious side effects, such as nausea, dyspepsia, fatigue, and metabolic acidosis, which side effects can affect patient compliance and/or necessitate the termination of treatment. Beta-blockers have increasingly become associated with serious pulmonary side effects

attributable to their effects on beta-2 receptors in pulmonary tissue. Sympathomimetics may cause tachycardia, arrhythmia and hypertension. Recently, certain prostaglandins and prostaglandin derivatives have been described in the art as being useful in reducing intraocular pressure. Typically, however, prostaglandin therapy for the treatment of elevated intraocular pressure is attended by undesirable side-effects, such as irritation and hyperemia of varying severity and duration. There is therefore a continuing need for therapies which control elevated intraocular pressure associated with glaucoma without the degree of undesirable side-effects attendant to most conventional therapies.

Prostaglandins are metabolite derivatives of arachidonic acid. Arachidonic acid in the body is converted to prostaglandin  $G_2$ , which is subsequently converted to prostaglandin  $H_2$ . Other naturally occurring prostaglandins are derivatives of prostaglandin  $H_2$ . A number of different types of prostaglandins have been discovered including A, B, D, E, F, G, I and J-Series prostaglandins (EP 0 561 073 A1). Of interest in the present invention are compounds which are believed to exhibit IOP lowering mechanisms similar to those exhibited by  $PGD_2$  (a D-series prostaglandin of formula I) and  $PGF_{2\alpha}$  (an F-series prostaglandin of formula II):



The relationship between  $PGD_2$  receptor activation and IOP lowering effects is not well understood. Various publications have reported that  $PGD_2$  receptor activation leads to second messenger activation and in particular, to the stimulation of adenylate cyclase and resultant increases in cAMP levels (Thierauch, *Prostaglandins and their Receptors: II. Receptor Structure and Signal Transduction*, Journal of Hypertension, volume 12, pages 1-5 (1994). Regardless of the mechanism,  $PGD_2$  has been shown to lower IOP (Nakajima,

*Effects of Prostaglandin D<sub>2</sub> and its analogue, BW245C, on Intraocular Pressure in Humans*, Graefe's Archive Ophthalmology, volume 229, pages 411-413 (1991)). Thus, it has been of interest in the ophthalmic field to develop synthetic PGD<sub>2</sub> analogs with IOP lowering efficacy.

5

Synthetic PGD<sub>2</sub>-type analogs have been pursued in the art (Graefe's Archive Ophthalmology, volume 229, pages 411-413 (1991)). Though PGD<sub>2</sub>-type molecules lower IOP, these types of molecules have also been associated with undesirable side effects resulting from topical ophthalmic dosing. Such effects have included an initial increase in IOP, conjunctival hyperemia, increases in microvascular permeability, and increases in eosinophile infiltration (Alm, *The Potential of Prostaglandin Derivatives in Glaucoma Therapy*, Current Opinion in Ophthalmology, volume 4, No. 11, pages 44-50 (1993)).

Similarly, the relationship of PGF<sub>2 $\alpha$</sub>  receptor activation and IOP lowering effects is not well understood. It is believed that PGF<sub>2 $\alpha$</sub>  receptor activation leads to increased outflow of aqueous humor. Regardless of the mechanism, PGF<sub>2 $\alpha$</sub>  and certain of its analogs have been shown to lower IOP (Giuffre, *The Effects of Prostaglandin F<sub>2 $\alpha$</sub>  the Human Eye*, Graefe's Archive Ophthalmology, volume 222, pages 139-141 (1985); and Kerstetter et al., *Prostaglandin F<sub>2 $\alpha$</sub> -1-Isopropylester Lowers Intraocular Pressure Without Decreasing Aqueous Humor Flow*, American Journal of Ophthalmology, volume 105, pages 30-34 (1988)). Thus, it has been of interest in the field to develop synthetic PGF<sub>2 $\alpha$</sub>  analogs with IOP lowering efficacy.

Synthetic PGF<sub>2 $\alpha$</sub> -type analogs have been pursued in the art (Graefe's Archive Ophthalmology, volume 229, pages 411-413 (1991)). Though PGF<sub>2 $\alpha$</sub> -type molecules lower IOP, a number of these types of molecules have also been associated with undesirable side effects resulting from topical ophthalmic dosing. Such effects include an initial increase in IOP, breakdown of the blood aqueous barrier and conjunctival

hyperemia (Alm, *The Potential of Prostaglandin Derivatives in Glaucoma Therapy*, Current Opinion in Ophthalmology, volume 4, No. 11, pages 44-50 (1993)).

Based on the foregoing, a need exists for the development of molecules that may activate the PGD<sub>2</sub> and/or PGF<sub>2α</sub> receptors, yielding a more efficacious lowering of IOP, while exhibiting fewer or reduced side effects.

An agent which exhibits the same or improved efficacy, but with reduced side effects when compared to other agents, is said to have an improved therapeutic profile. It is an object of this invention to provide a class of IOP lowering agents with an improved therapeutic profile over their PGF<sub>2α</sub> and PGD<sub>2</sub> counterparts, and methods of their use. It has now unexpectedly been discovered that the presently claimed *cis*-Δ<sup>4</sup> analogs of PGF<sub>2α</sub> and PGD<sub>2</sub> meet this objective. Certain *cis*-Δ<sup>4</sup> analogs of PGF<sub>2α</sub> (Nedy and Johnson, J. Org. Chem., 45:6, 1121 (1980); Bowler *et. al.* Prostaglandins, 17:6, 789 (1979); DE 2,716,972; DE 2,637,384; DE 2,623,139; US 3,954,835) and PGD<sub>2</sub> (EP0 299,914 B1) are known in the art. The metabolic profiles of *cis*-Δ<sup>4</sup> PGF<sub>2α</sub> and (15*S*)-15-methyl- *cis*-Δ<sup>4</sup> PGF<sub>2α</sub> relative to the corresponding *cis*-Δ<sup>5</sup> isomers have also been discussed in the art (Green *et. al.*, Eur. J. Biochem., 62, 527 (1976); Green, Prostaglandins, 15:5, 813 (1978); Hansson, Prostaglandins, 18:5, 745 (1979); Tarpley and Sun, J. Med. Chem., 21:3, 288 (1978)). However, the surprisingly enhanced therapeutic profiles of such compounds and the novel compounds of the present invention in the treatment of glaucoma are neither disclosed nor suggested in that art.


## Summary of the Invention

The present invention is directed to compositions and methods of their use in treating IOP and ocular hypertension. In particular, the present invention provides certain classes of *cis*-Δ<sup>4</sup> prostaglandin analogs, and methods of their use in treating glaucoma and ocular hypertension. As previously stated, the mechanism of action by which PGD<sub>2</sub> and PGF<sub>2α</sub> type prostaglandins lower IOP is not well understood. Nevertheless, without being

bound by any theories, the inventors postulate that the compounds of the present invention, with a *cis* double bond between carbons 4 and 5 in the alpha ( $\alpha$ ) chain, are less susceptible to metabolic degradation than are their non-*cis*- $\Delta^4$  counterparts, and therefore enjoy a longer *in vivo* half-life than the non- *cis*- $\Delta^4$  compounds. A longer *in vivo* half life should enable relatively lower dosing and a reduction in side effects.

### Detailed Description of the Invention

It has unexpectedly been found that *cis*- $\Delta^4$  analogs of the present invention exhibit an improved therapeutic profile in the treatment of glaucoma and ocular hypertension when compared to natural prostaglandins and many of their known analogs. The *cis*- $\Delta^4$  analogs of the present invention are heptenoic acid derivatives having the following formula **III**:




wherein:

A = CO<sub>2</sub>R, CONR<sup>1</sup>R<sup>2</sup>, CH<sub>2</sub>OR<sup>3</sup>, or CH<sub>2</sub>NR<sup>4</sup>R<sup>5</sup>; where R = H or cationic salt moiety, or CO<sub>2</sub>R = pharmaceutically acceptable ester moiety; R<sup>1</sup>, R<sup>2</sup> = same or different = H or alkyl; R<sup>3</sup> = H, acyl, or alkyl; R<sup>4</sup>, R<sup>5</sup> = same or different = H, acyl, or alkyl, with the proviso that if one of R<sup>4</sup>, R<sup>5</sup> = acyl, then the other = H or alkyl;

n = 0 or 2;

L = OR<sup>6</sup> in the  $\alpha$  configuration, where R<sup>6</sup> = H, alkyl, or acyl; or L = halo in either configuration;




---- = single or *trans* double bond;

$D, D^1$ , taken together =  $\text{OCH}_2\text{CH}_2\text{O}$ ; or  $D, D^1 = \text{different} = \text{H and OR}^8$ , where  $R^8 = \text{H, alkyl, or acyl}$ ; or  $D = \text{fluorine in the } \alpha \text{ configuration, and } D^1 = \text{H in the } \beta \text{ configuration}$ ;

$X = (\text{CH}_2)_m \text{ or } (\text{CH}_2)_m\text{O}$ , where  $m = 1-6$ ; and

$Y = \text{a phenyl ring optionally substituted with alkyl, halo, trihalomethyl, alkoxy, acyl, acyloxy, amino, alkylamino, acylamino, or hydroxy}$ ; or

$X-Y = (\text{CH}_2)_p Y^1$ ; where  $p = 0-6$ ; and



wherein:

$W = \text{CH}_2, \text{O}, \text{S(O)}_q, \text{NR}^9, \text{CH}_2\text{CH}_2, \text{CH=CH}, \text{CH}_2\text{O}, \text{CH}_2\text{S(O)}_q, \text{CH=N, or } \text{CH}_2\text{NR}^9$ ; where  $q = 0-2$ , and  $R^9 = \text{H, alkyl, or acyl}$ ;

$Z = \text{H, alkyl, alkoxy, acyl, acyloxy, halo, trihalomethyl, amino, alkylamino, acylamino, or hydroxy}$ ; and

---- = single or double bond;




or X-Y = cyclohexyl;

with the proviso that the following compounds of formula **III** be excluded: those wherein:

L = OR<sup>6</sup> in the  $\alpha$  configuration, where R<sup>6</sup> is as defined above;

5



--- = *trans* double bond;

10 D, D<sup>1</sup> = different = H and OR<sup>8</sup>, where R<sup>8</sup> is as defined above;

X = CH<sub>2</sub>CH<sub>2</sub> or CH<sub>2</sub>O; and

Y = a phenyl ring, optionally substituted with halo.

1  
15


For purposes of the foregoing and following definitions, the term “pharmaceutically acceptable ester” means any ester that would be suitable for therapeutic administration to a patient by any conventional means without significant deleterious health consequences; and “ophthalmically acceptable ester” means any pharmaceutically acceptable ester that would be suitable for ophthalmic application, i.e. non-toxic and non-irritating. Preferred are alkyl esters. Most preferred are C<sub>2</sub>-C<sub>4</sub> alkyl esters, and especially isopropyl esters. For compounds of formula **III** where L = halo, preferred is chlorine in the  $\beta$  configuration.

25 Preferred for use in the methods and compositions of the present invention are those compounds of formula **III** above, wherein:

A = CO<sub>2</sub>R, where R = H; or CO<sub>2</sub>R = pharmaceutically acceptable ester moiety selected from the group consisting of ophthalmically acceptable ester moieties, where R = alkyl;

n = 0;

L = OH in the  $\alpha$  configuration;



5


D, D<sup>1</sup>, taken together = OCH<sub>2</sub>CH<sub>2</sub>O; or D = OH in the  $\alpha$  configuration, and D<sup>1</sup> = H in the  $\beta$  configuration; or D = fluorine in the  $\alpha$  configuration, and D<sup>1</sup> = H in the  $\beta$  configuration;

X = CH<sub>2</sub>CH<sub>2</sub> or CH<sub>2</sub>O; and

10

Y = phenyl, optionally substituted with halo or trihalomethyl; or

X-Y =




15

with the proviso that the following compounds of formula III be excluded, wherein:

A = CO<sub>2</sub>R, where R = H or alkyl;

20 n = 0;

L = OH in the  $\alpha$  configuration;



25

--- = *trans* double bond;

D = OH in the  $\alpha$  configuration, and D<sup>1</sup> = H in the  $\beta$  configuration;

X = CH<sub>2</sub>CH<sub>2</sub> or CH<sub>2</sub>O; and

5

Y = a phenyl ring, optionally substituted with halo.

Especially preferred are the following compounds:

Compound Number	Compound Name	Compound Structure
<b>IV</b>	(4Z,13E)-(9S,11R,15R)-16-(3-Trifluoromethylphenoxy)-9,11,15-trihydroxy-17,18,19,20-tetranor-4,13-prostadienoic acid isopropyl ester	
<b>V</b>	(4Z,13E)-(9S,11R,15S)-15-(2-Indanyl)-9,11,15-trihydroxy-16,17,18,19,20-pentanor-4,13-prostadienoic acid isopropyl ester	
<b>VI</b>	(4Z,13E)-(9S,11R)-16-(3-Chlorophenoxy)-9,11-dihydroxy-15-(1,3-dioxolan-2-yl)-17,18,19,20-tetranor-4,13-prostadienoic acid isopropyl ester	
<b>VII</b>	(4Z,13E)-(9S,11S,15R)-9,15-Dihydroxy-11-fluoro-16-[(3-trifluoromethyl)phenoxy]-17,18,19,20-tetranor-4,13-prostadienoic acid isopropyl ester	
<b>VIII</b>	Isopropyl [2R(1E,3R),3S(4Z),4R]-7-[Tetrahydro-2-[4-(3-chlorophenoxy)-3-hydroxy-1-butenyl]-4-hydroxy-3-furanyl]-4-heptenoate	
<b>IX</b>	(4Z)-(9S,11R,15R)-16-(3-Chlorophenoxy)-9,11-dihydroxy-15-fluoro-17,18,19,20-tetranor-4-prostenoic acid isopropyl ester	

10


Included within the scope of the present invention are the individual enantiomers of the title compounds, as well as their racemic and non-racemic mixtures. The individual enantiomers can be enantioselectively synthesized from the appropriate enantiomerically pure or enriched starting material by means such as those described below. Alternatively, they may be enantioselectively synthesized from racemic/non-racemic or achiral starting materials. (*Asymmetric Synthesis* by J. D. Morrison and J. W. Scott, Eds., Academic Press Publishers: New York, 1983-1985 (five volumes) and *Principles of Asymmetric Synthesis* by R.E. Gawley and J. Aube, Eds., Elsevier Publishers: Amsterdam, 1996). They may also be isolated from racemic and non-racemic mixtures by a number of known methods, e.g. by purification of a sample by chiral HPLC (*A Practical Guide to Chiral Separations by HPLC*, G. Subramanian, Ed., VCH Publishers: New York, 1994; *Chiral Separations by HPLC*, A.M. Krstulovic, Ed., Ellis Horwood Ltd. Publishers, 1989), or by enantioselective hydrolysis of a carboxylic acid ester sample by an enzyme (Ohno, M.; Otsuka, M. Organic Reactions, volume 37, page 1 (1989)). Those skilled in the art will appreciate that racemic and non-racemic mixtures may be obtained by several means, including without limitation, nonenantioselective synthesis, partial resolution or even mixing samples having different enantiomeric ratios.

Believed to be novel are compounds of formula **III** which contain a bicyclic aromatic group on the terminus of the  $\omega$  chain, *i.e.*, those wherein:

A = CO<sub>2</sub>R, CONR<sup>1</sup>R<sup>2</sup>, CH<sub>2</sub>OR<sup>3</sup>, or CH<sub>2</sub>NR<sup>4</sup>R<sup>5</sup>; where R = H or cationic salt moiety, or CO<sub>2</sub>R forms a pharmaceutically acceptable ester moiety; R<sup>1</sup>, R<sup>2</sup> = same or different = H or alkyl; R<sup>3</sup> = H, acyl, or alkyl; R<sup>4</sup>, R<sup>5</sup> = same or different = H, acyl, or alkyl, with the proviso that if one of R<sup>4</sup>, R<sup>5</sup> = acyl, then the other = H or alkyl;


n = 0 or 2;

L = OR<sup>6</sup> in the  $\alpha$  configuration, where R<sup>6</sup> = H, alkyl, or acyl;



---- = single or *trans* double bond;

- 5 D, D', taken together = OCH<sub>2</sub>CH<sub>2</sub>O; or D, D' = different = H and OR<sup>8</sup>, where R<sup>8</sup> = H, alkyl, acyl; or D = fluorine in the α configuration, and D' = H in the β configuration;

$$X-Y = (CH_2)_p Y^1; \text{ where } p = 0-6; \text{ and}$$


wherein:

W = CH<sub>2</sub>, O, S(O)<sub>q</sub>, NR<sup>9</sup>, CH<sub>2</sub>CH<sub>2</sub>, CH=CH, CH<sub>2</sub>O, CH<sub>2</sub>S(O)<sub>q</sub>, CH=N, or CH<sub>2</sub>NR<sup>9</sup>; where q = 0-2, and R<sup>9</sup> = H, alkyl, or acyl;

Z = H, alkyl, alkoxy, acyl, acyloxy, halo, trihalomethyl, amino, alkylamino, acylamino, or hydroxy; and

---- = single or double bond.

Other related compounds within the scope of the present invention are known and their syntheses are either described in the literature or can be achieved by methods similar to those described in the literature or otherwise known to those of skill in the art. . See, for example, J. Org. Chem., 45:6, 1121 (1980); Prostaglandins, 17(6):789 (1979); Eur. J. Biochem., 62:527 (1976); DE 2,826,462; DE 2,716,972; DE 2,623,139; EPO 299914 B1. The contents of these references are by this reference incorporated herein in their entirety.


In the foregoing illustrations, as well as those provided hereinafter, wavy line attachments indicate either the alpha ( $\alpha$ ) or beta ( $\beta$ ) configuration. The carbon numbering is as indicated in structural formula **III** (even when  $n = 2$ ). A hatched line, as used e.g. at carbon 9, indicates the  $\alpha$  configuration. A solid triangular line, as used e.g. at carbon 12, indicates the  $\beta$  configuration. Dashed lines on bonds, e.g. between carbons 13 and 14, indicate a single or double bond. Two solid lines between carbons indicate a double bond of the specified configuration.

In the following Examples 1-6, the following standard abbreviations are used:

g = grams (mg = milligrams); mol = moles (mmol = millimoles); mL = milliliters; mm Hg = millimeters of mercury; mp = melting point; bp = boiling point; h = hours; and min = minutes. In addition, "NMR" refers to nuclear magnetic resonance spectroscopy and "MS" refers to mass spectrometry.

## EXAMPLE 1:

## Synthesis of IV




(4Z,13E)-(9S,11R,15R)-16-(3-Trifluoromethylphenoxy)-9,11,15-trihydroxy-17,18,19,20-tetranor-4,13-prostadienoic acid isopropyl ester (IV)

Reaction of diol 1 (for preparation, see U.S. Patent Number 4,321,275, which is  
5 incorporated by this reference) with 3,4-dihydro-2H-pyran in CH<sub>2</sub>Cl<sub>2</sub> at 0 °C in the  
presence of *p*-toluenesulfonic acid (TsOH) affords THP ether 2, which is reduced with  
diisobutylaluminum hydride (DIBAL-H) in toluene at -78 °C to afford lactol 3. Wittig  
reaction of 3 with Ph<sub>3</sub>P<sup>+</sup>CH<sub>2</sub>OCH<sub>3</sub> Cl<sup>-</sup> in THF at 0 °C in the presence of potassium *t*-  
butoxide (KOBU<sup>t</sup>) provides enol ether 4, which is hydrolyzed with TsOH in THF/water to  
10 yield lactol 5. Wittig reaction of 5 with Ph<sub>3</sub>P<sup>+</sup>(CH<sub>2</sub>)<sub>3</sub>CO<sub>2</sub>H Br<sup>-</sup> in THF at 0 °C in the  
presence of KOBU<sup>t</sup>, followed by alkylation of the product carboxylic acid with isopropyl  
iodide in acetone in the presence of DBU, yields IV after purification *via* silica gel  
chromatography.



## EXAMPLE 2:

## Synthesis of V




(4Z,13E)-(9S,11R,15S)-15-(2-Indanyl)-9,11,15-trihydroxy-16,17,18,19,20-pentanoic acid isopropyl ester (V)

Reaction of diol **6** (for preparation, see U.S. Patent Number 4,152,527, which is  
5 incorporated by this reference) with 3,4-dihydro-2*H*-pyran in CH<sub>2</sub>Cl<sub>2</sub> at 0 °C in the  
presence of TsOH affords THP ether **7**, which is reduced with diisobutylaluminum hydride  
DIBAL-H in toluene at -78 °C to afford lactol **8**. Wittig reaction of **8** with  
Ph<sub>3</sub>P<sup>+</sup>CH<sub>2</sub>OCH<sub>3</sub> Cl<sup>-</sup> in THF at 0 °C in the presence KOBu<sup>t</sup> provides enol ether **9**, which is  
hydrolyzed with TsOH in THF/water to yield lactol **10**. Wittig reaction of **10** with  
10 Ph<sub>3</sub>P<sup>+</sup>(CH<sub>2</sub>)<sub>3</sub>CO<sub>2</sub>H Br<sup>-</sup> in THF at 0 °C in the presence of KOBu<sup>t</sup>, followed by alkylation of  
the product carboxylic acid with isopropyl iodide in acetone in the presence of DBU,  
yields **V** after purification *via* silica gel chromatography.

## EXAMPLE 3:

## Synthesis of VI




(4Z,13E)-(9S,11R)-16-(3-Chlorophenoxy)-9,11-dihydroxy-15-(1,3-dioxolan-2-yl)-  
17,18,19,20-tetranor-4,13-prostadienoic acid isopropyl ester (VI)

Ketalization of enone **11** (for preparation, see published European Patent

- 5 Application No. EP 639563 A2, which is incorporated by this reference) with  
(Me<sub>3</sub>SiOCH<sub>2</sub>)<sub>2</sub> in CH<sub>2</sub>Cl<sub>2</sub> at -78 °C in the presence of Me<sub>3</sub>SiOSO<sub>2</sub>CF<sub>3</sub> affords ketal **12**,  
which is debenzoylated with potassium carbonate in methanol to provide alcohol **13**.  
Reaction of **13** with 3,4-dihydro-2H-pyran in CH<sub>2</sub>Cl<sub>2</sub> at 0 °C in the presence TsOH affords  
**14**, reduction of which with DIBAL-H in toluene at -78 °C provides lactol **15**. Wittig  
10 reaction of lactol **15** with Ph<sub>3</sub>P<sup>+</sup>CH<sub>2</sub>OMe Cl<sup>-</sup> in the the presence of KOBu<sup>t</sup> in THF affords  
enol ether **16** as a mixture of enol ether olefin geometrical isomers. Treatment of this  
mixture with TsOH in THF/water affords lactol **17**. Wittig reaction of **17** with  
Ph<sub>3</sub>P<sup>+</sup>(CH<sub>2</sub>)<sub>3</sub>CO<sub>2</sub>H Br<sup>-</sup> in the presence of KOBu<sup>t</sup> in THF, followed by treatment of an  
acetone solution of the resulting carboxylic acid with DBU and isopropyl iodide, yields **VI**  
15 after purification *via* silica gel chromatography.

## EXAMPLE 4:

## Synthesis of VII



(4Z,13E)-(9S,11S,15R)-9,15-Dihydroxy-11-fluoro-16-[(3-trifluoromethyl)phenoxy]-  
17,18,19,20-tetranor-4,13-prostadienoic acid isopropyl ester (VII)

Reduction of aldehyde **18** with NaBH<sub>4</sub> in methanol/CH<sub>2</sub>Cl<sub>2</sub> at 0 °C affords alcohol

5 **19**, which is treated with Ph<sub>2</sub>Bu<sup>t</sup>SiCl in CH<sub>2</sub>Cl<sub>2</sub> in the presence of imidazole and 4-(dimethylamino)pyridine to afford silyl ether **20**. Debenzoylation is effected using potassium carbonate in methanol to provide alcohol **21**, which is treated with (diethylamino)sulfur trifluoride (DAST) in CH<sub>2</sub>Cl<sub>2</sub> at 0 °C. After work-up and purification of the reaction *via* silica gel chromatography, the residue is stirred overnight


10 with catalytic OsO<sub>4</sub> in acetone in the presence of stoichiometric *N*-methylmorpholine *N*-oxide to provide fluoride **22** after purification *via* silica gel chromatography. Reduction of **22** with DIBAL-H affords lactol **23**, which is condensed with Ph<sub>3</sub>P<sup>+</sup>CH<sub>2</sub>OMe Cl<sup>-</sup> in the the presence of KOBu<sup>t</sup> in THF to yield enol ether **24**. Acidic hydrolysis using TsOH in THF/water gives lactol **25**, which is reacted with Ph<sub>3</sub>P<sup>+</sup>(CH<sub>2</sub>)<sub>3</sub>CO<sub>2</sub>H Br<sup>-</sup> in the presence of

15 KOBu<sup>t</sup> in THF, followed by treatment of an acetone solution of the resulting carboxylic acid with DBU and isopropyl iodide, to afford alkene **26**. Treatment of **26** with 3,4-dihydro-2*H*-pyran in CH<sub>2</sub>Cl<sub>2</sub> at 0 °C in the presence TsOH provides THP ether **27**, which is desilylated using tetra-*n*-butylammonium fluoride in THF to give alcohol **28**. Swern oxidation of **28** (oxalyl chloride, DMSO, CH<sub>2</sub>Cl<sub>2</sub>, NEt<sub>3</sub>, -78 °C) affords aldehyde **29**,

20 which is treated with (MeO)<sub>2</sub>P(O)CH<sub>2</sub>C(O)CH<sub>2</sub>OC<sub>6</sub>H<sub>4</sub>-*m*-CF<sub>3</sub>, NEt<sub>3</sub>, and LiCl in THF to provide enone **30**. Luche reduction of **30** (NaBH<sub>4</sub>, CeCl<sub>3</sub>, MeOH, 0 °C) gives allyl alcohol **31** as a mixture of C-15 diastereomers. Acidic deprotection of **31** using aqueous HCl in isopropanol, followed by purification *via* silica gel chromatography, gives **VII**.

## EXAMPLE 5:

## Synthesis of VIII



(3aR,4S,6aR)-4-(*tert*-Butyldiphenylsilyloxy)methylhexahydrofuro[3,4-b]furan-2-one (32)

A mixture of alcohol **31** (for preparation, see U.S. Patent No. 4,133,948, which is incorporated by this reference) (5.0 g, 31.6 mmol) and imidazole (4.3 g, 63.2 mmol) was dissolved in 100 mL of anhydrous DMF. To this solution *tert*-butyldiphenylsilyl chloride (10.4 g, 38.0 mmol) was added and the resulting mixture was stirred at room temperature for 14 h. The solvent was evaporated and the residue was taken up in 100 mL of EtOAc, washed with water (2X50 mL), dilute aqueous solution of HCl (2X50 mL) and brine and dried (MgSO<sub>4</sub>). The solvent was evaporated and the crude was purified by chromatography on silica gel to afford **32** (12.4 g, quantitative yield) as a white solid: *R<sub>f</sub>* 0.6 (60% EtOAc/hexanes). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 7.65 (m, 4H), 7.42 (m, 6H), 5.10 (m, 1H), 4.25 (dd, *J* = 12, 4 Hz, 1H), 4.05 (dd, *J* = 12, 2 Hz, 1H), 3.85 (m, 1H), 3.75 (m, 2H), 3.00 (m, 1H), 2.82 (dd, *J* = 16, 7 Hz, 1H), 2.45 (dd, *J* = 16, 2 Hz, 1H), 1.05 (s, 9H).

Isopropyl [2S,3S(4Z),4R]-7-[Tetrahydro-2-(*tert*-butyldiphenylsilyloxy)methyl-4-hydroxy-3-furanyl]-4-heptenoate (35)

A solution of the lactone **32** (5.7 g, 14.5 mmol) in 150 mL of anhydrous THF was cooled to -78 °C under an inert atmosphere, and to it DIBAL-H (14.5 mL, 1.5 M in toluene, 21.7 mmol) was added dropwise. The resulting mixture was stirred at -78 °C for 1.5 h and was then quenched at the same temperature by the addition of 5 mL of methanol. The reaction was warmed to room temperature, an equal volume of a saturated aqueous solution of potassium sodium tartrate was added to it, and the resulting slurry was stirred at room temperature for 1 h. The layers were separated, and the aqueous layer was extracted with 3X25 mL of EtOAc. The organic layers were combined and washed with brine and dried (MgSO<sub>4</sub>). The solution was filtered and concentrated and the crude was purified by passage through a short column of silica gel to afford the intermediate lactol (5.6 g, quantitative yield) as a colorless oil: *R<sub>f</sub>* 0.5 (60% EtOAc/hexanes).

A suspension of (methoxymethyl)triphenylphosphonium chloride (2.5 g, 7.5 mmol) in 70 mL of dry THF was cooled to 0 °C under a N<sub>2</sub> atmosphere. To this solution



potassium *tert*-butoxide (*t*-BuOK, 9.0 mL, 1.0 M in THF, 9.0 mmol) was added dropwise, and stirring was continued at 0 °C for an additional 20 min. At this time a solution of the lactol obtained above (1.0 g, 2.5 mmol) in 30 mL of dry THF was added to it, and the resulting mixture was stirred at 0 °C for 1.5 h. The reaction was then worked up by  
5 pouring it into 50 mL of a saturated aqueous solution of KH<sub>2</sub>PO<sub>4</sub>, the layers were separated and aqueous layer was extracted with 3X25 mL of EtOAc. The combined organic layers were washed with water and brine, and dried (MgSO<sub>4</sub>); solvent removal and chromatography of the crude on silica afforded the enolether **33** (0.89 g, 83% yield) as a colorless liquid: R<sub>f</sub> 0.6 (60% EtOAc/hexanes).

10 A solution containing enolether **33** (2.45 g, 5.7 mmol), *p*-toluenesulfonic acid (0.1 g) and water (10 mL) in 150 mL of THF was heated at reflux for 3 h. The mixture was then cooled to room temperature and poured into 50 mL of a saturated aqueous solution of NaHCO<sub>3</sub>. The layers were separated and aqueous layer was extracted with EtOAc. The  
15 organic extracts were combined and dried (MgSO<sub>4</sub>) and the crude product was subjected to chromatography on silica to afford **34** (1.44 g, 60% yield) as a colorless liquid. This material was used in the next reaction: R<sub>f</sub> 0.28 (50% EtOAc/hexanes).

A suspension of (3-carboxypropyl)triphenylphosphonium bromide (4.5 g, 10.5  
20 mmol) in 70 mL of dry THF was cooled to 0 °C and to it *t*-BuOK (21.0 mL, 1.0 M in THF, 21.0 mmol) was added dropwise. The resulting solution was stirred for 30 min at 0 °C and to it a solution of the lactol **34** (1.44 g, 3.5 mmol) in 30 mL of dry THF was added dropwise over a period of 10 min. The reaction was allowed to warm to room temperature gradually, and was stirred at that temperature for 14 h. The mixture was then poured into  
25 50 mL of a saturated aqueous solution of KH<sub>2</sub>PO<sub>4</sub>, and extracted with 3X25 mL of EtOAc. The organic extracts were combined and washed with brine, and dried (MgSO<sub>4</sub>). Filtration and solvent removal afforded an oily residue which was used immediately in the subsequent step.

30 The crude product from above was dissolved in 40 mL of acetone and the solution was treated with DBU (12.0 mL, 84 mmol) at room temperature for 10 min. Isopropyl

iodide (7.0 mL, 70 mmol) was then introduced and the resulting mixture was stirred at room temperature for 18 h. Solvent was evaporated, and the residue was dissolved in 50 mL of EtOAc. This solution was washed sequentially with 3X25 mL of a saturated aqueous solution of  $\text{KH}_2\text{PO}_4$ , 1X10 mL of water and brine and dried over anhydrous  $\text{MgSO}_4$ . Filtration, solvent removal and chromatography of the crude on silica gel afforded the desired isopropyl ester **35** (1.18 g, 65% yield from **34**) as a slightly yellow liquid:  $R_f$  0.2 (30% EtOAc/hexanes);  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  7.71 (m, 4H), 7.40 (m, 6H), 5.38 (m, 2H), 5.00 (septet,  $J = 6.4$  Hz, 1H), 4.38 (m, 1H), 3.65-4.00 (broad m, 5H), 1.90-2.50 (broad m, 7H), 1.55 (m, 2H), 1.23 (d,  $J = 7.2$  Hz, 6H), 1.05 (s, 9H); MS  $m/z$  at 547 for  $(\text{M}+\text{Na})^+$ .

Isopropyl [2S,3R(4Z),4R]-7-[Tetrahydro-2-hydroxymethyl-4-(tetrahydropyran-2-yl)oxy-3-furanyl]-4-heptenoate (**37**)

A solution of the alcohol **35** (1.18 g, 2.3 mmol) and 3,4-dihydro-2H-pyran (0.3 mL, 3.4 mmol) in 50 mL of  $\text{CH}_2\text{Cl}_2$  was cooled to  $0^\circ\text{C}$  and to it a catalytic amount of p-toluenesulfonic acid (10 mg) was added. The resulting mixture was stirred at  $0^\circ\text{C}$  for 25 min and was then quenched by the addition of 25 mL of a saturated aqueous solution of  $\text{NaHCO}_3$ . The mixture was warmed to room temperature, the layers were separated and the aqueous layer was extracted with 3X25 mL of  $\text{CH}_2\text{Cl}_2$ . The organic layers were combined and washed with brine and dried ( $\text{K}_2\text{CO}_3$ ). The crude obtained after filtration and solvent removal was purified by passage through a short plug of silica to afford the intermediate tetrahydropyranyl ether **36** as colorless liquid:  $R_f$  0.4 (30% EtOAc/hexanes).

The silyl ether **36** thus obtained was dissolved in 20 mL of THF and the solution was treated with tetra-n-butylammonium fluoride (7.0 mL, 1.0 M in THF, 7.0 mmol) at room temperature for 2 h. The reaction mixture was then poured into water and was extracted with EtOAc (3X25 mL). The organic extracts were combined and dried ( $\text{MgSO}_4$ ), filtered and concentrated. The crude was subjected to chromatography on silica to afford the alcohol **37** (0.72 g, 85% yield from **35**) as a colorless liquid:  $R_f$  0.16 (50%

EtOAc/hexanes);  $^1\text{H-NMR}$  ( $d_6$ -DMSO)  $\delta$  (partial spectrum) 5.36 (m, 2H), 4.87 (septet,  $J = 6.5$  Hz, 1H), 4.60 (m, 2H), 1.18 (d,  $J = 7.2$  Hz, 6H).

Isopropyl [2S,3R(4Z),4R]-7-[Tetrahydro-2-formyl]-4-(tetrahydropyran-2-yl)oxy-3-furanyl]-4-heptenoate (38)

A solution of oxalyl chloride (2.0 mL, 2.0 M in  $\text{CH}_2\text{Cl}_2$ , 4.0 mmol) in 10 mL of dry  $\text{CH}_2\text{Cl}_2$  was cooled to  $-78^\circ\text{C}$ , and to it a solution of DMSO (0.56 mL, 8.0 mmol) in 5 mL of  $\text{CH}_2\text{Cl}_2$  was introduced dropwise. After the mixture was stirred for 3 min at  $-78^\circ\text{C}$ , a solution of the substrate **37** (0.72 g, 2.0 mmol) in 25 mL of  $\text{CH}_2\text{Cl}_2$  was added to it dropwise. The mixture was stirred for 15 min, at which time triethylamine (1.7 mL, 12.0 mmol) was introduced, and stirring was continued for an additional 15 min. The reaction was gradually warmed to room temperature and then poured into 50 mL of water. The layers were separated and the water layer was extracted with 3X25 mL of  $\text{CH}_2\text{Cl}_2$ . The combined organic extracts were washed with water and brine, and dried ( $\text{MgSO}_4$ ). Filtration and solvent removal, followed by chromatography of the crude on silica afforded the aldehyde **38** (0.69 g, 94% yield) as a pale yellow liquid:  $R_f$  0.3 (50% EtOAc/hexanes);  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  (partial spectrum) 9.66 (d,  $J = 3$  Hz, 1H), 5.37 (m, 2H), 5.0 (septet,  $J = 6.5$  Hz, 1H), 1.24 (d,  $J = 7.2$  Hz, 6H).

Isopropyl [2R(1E),3R(4Z),4R]-7-[Tetrahydro-2-[4-(3-chlorophenoxy)-3-oxo-1-butenyl]-4-(tetrahydropyran-2-yl)oxy-3-furanyl]-4-heptenoate (39)

A mixture of the aldehyde **38** (0.32 g, 0.87 mmol), dimethyl-3-(3-chlorophenoxy)-2-oxopropylphosphonate (1.0 g, 3.5 mmol) and  $\text{LiCl}$  (0.15 g, 3.5 mmol) was taken up in 40 mL of dry THF, and the solution was cooled to  $0^\circ\text{C}$  under a  $\text{N}_2$  atmosphere. To this solution triethylamine (0.5 mL, 3.5 mmol) was added dropwise, and the resulting slurry was stirred at  $0^\circ\text{C}$  for 1 h. The reaction was then quenched by pouring it into 50 mL of a saturated aqueous solution of  $\text{KH}_2\text{PO}_4$ . The organic layer was separated and the aqueous layer was extracted with 3X25 mL of EtOAc. The organic extracts were combined and washed with water and brine and dried ( $\text{MgSO}_4$ ). The crude product mixture was

subjected to chromatography on silica to afford the enone **39** (0.34 g, 73% yield) as a pale yellow liquid:  $R_f$  0.6 (60% EtOAc/hexanes);  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  (partial spectrum) 6.70-7.20 (broad m, 5H), 6.12 (d,  $J = 16.7$  Hz, 1H), 5.36 (m, 2H), 5.0 (septet,  $J = 6.5$  Hz, 1H), 4.73 (s, 2H), 1.23 (d,  $J = 7.5$  Hz, 6H).

Isopropyl [2R(1E,3RS),3R(4Z),4R]-7-[Tetrahydro-2-[4-(3-chlorophenoxy)-3-hydroxy-1-butenyl]-4-(tetrahydropyran-2-yl)oxy-3-furanyl]-4-heptenoate (**40**)

A mixture of the enone **39** (0.34 g, 0.64 mmol) and  $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$  (0.47 g, 1.27 mmol) was dissolved in 30 mL of methanol and the solution was cooled to  $-5^\circ\text{C}$ .  $\text{NaBH}_4$  (47 mg, 1.27 mmol) was added to the solution in small portions over a period of 3 min. The mixture was stirred for an additional 3 min and the reaction was then quenched at  $-5^\circ\text{C}$  by the addition of 10 mL of a saturated aqueous solution of  $\text{NH}_4\text{Cl}$ . The resulting slurry was warmed to room temperature and partitioned between  $\text{CHCl}_3$  and water. The aqueous layer was extracted with 3X25 mL of  $\text{CHCl}_3$  and the combined organic extracts were washed with 2X10 mL of water and brine. The organic layer was dried, filtered and concentrated and the crude was purified by chromatography on silica to afford the reduction product **40** (0.30 g, 87% yield) as a colorless liquid:  $R_f$  0.24 (50% EtOAc/hexanes).


Isopropyl [2R(1E,3R),3S(4Z),4R]-7-[Tetrahydro-2-[4-(3-chlorophenoxy)-3-hydroxy-1-butenyl]-4-hydroxy-3-furanyl]-4-heptenoate (**VIII**)

The allyl alcohol **40** (0.30 g, 0.55 mmol) was dissolved in a mixture of 10 mL of methanol and 1.0 mL of water, and the solution was cooled to  $0^\circ\text{C}$ . Approximately 10 drops of 12  $N$   $\text{HCl}$  was added to it dropwise, and stirring was continued at  $0^\circ\text{C}$  for 15 min and then at room temperature for 1 h. The reaction was then quenched by the addition of solid  $\text{NaHCO}_3$ , and the suspension was partitioned between  $\text{CHCl}_3$  and water. The layers were separated and the aqueous layer was extracted with 3X25 mL of  $\text{CHCl}_3$ . The organic extracts were combined and washed with water (2X10 mL) and brine and dried ( $\text{Na}_2\text{SO}_4$ ). Filtration and solvent removal gave an oil which was subjected to silica gel

chromatography to separate the two diastereomers. **VIII** (61 mg, 25% yield) was obtained as colorless liquid:  $R_f$  0.15 (60% EtOAc/hexanes);  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  7.17 (m, 1H), 6.90 (m, 2H), 6.78 (m, 1H), 5.84 (m, 2H), 5.35 (m, 2H), 5.00 (septet,  $J = 6.4$  Hz, 1H), 4.55 (m, 1H), 4.40 (m, 1H), 3.80-4.15 (broad m, 5H), 1.90-2.65 (broad m, 8H), 1.75 (m, 2H), 1.45 (m, 2H), 1.21 (d,  $J = 7.4$  Hz, 6H);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  173.08, 159.19, 134.90, 132.69, 130.68, 130.57, 130.26, 128.07, 121.35, 115.09, 113.04, 82.21, 75.45, 72.62, 71.83, 70.12, 67.94, 50.84, 34.36, 25.78, 24.55, 22.70, 21.89, 21.80; HRMS  $m/z$  calculated for  $\text{C}_{24}\text{H}_{33}\text{O}_6\text{ClNa}$  ( $\text{M} + \text{Na}^+$ ) 475.185884, found 475.18588.

## EXAMPLE 6:

## Synthesis of IX



(4Z)-(9S,11R,15R)-16-(3-Chlorophenoxy)-9,11-dihydroxy-15-fluoro-17,18,19,20-tetranor-4-prostenoic acid isopropyl ester (IX)

Ketone reduction of 11 with (+)-*B*-chlorodiisopinocampheylborane in THF at 0 °C  
5 affords C-15  $\beta$ -hydroxy isomer 41 after chromatographic purification. Olefin reduction  
using Pd/C under a hydrogen atmosphere in ethyl acetate solvent provides alcohol 42,  
which is fluorinated using DAST to yield  $\alpha$ -fluoride 43. Debenzoylation with potassium  
carbonate in methanol gives alcohol 44, which is treated with 3,4-dihydro-2*H*-pyran in  
methylene chloride in the presence of TsOH to afford THP ether 45. Reduction of 45 to  
10 lactol 46 is effected with DIBAL-H at -78 °C in toluene, and Wittig condensation of 46  
with  $\text{Ph}_3\text{P}^+\text{CH}_2\text{OMe Cl}^-$  in the presence of  $\text{KOBU}^+$  in THF yields enol ether 47.  
Acidic hydrolysis using TsOH in THF and water gives lactol 48, which is reacted with  
 $\text{Ph}_3\text{P}^+(\text{CH}_2)_3\text{CO}_2\text{H Br}^-$  in the presence of  $\text{KOBU}^+$  in THF, followed by treatment of an  
acetone solution of the resulting carboxylic acid with DBU and isopropyl iodide, to afford  
15 **IX**.

The *cis*- $\Delta^4$  analogs of the present invention may be formulated in various  
pharmaceutical compositions for administering to humans and other mammals as a  
treatment of glaucoma or ocular hypertension. As used herein, the term “pharmaceutically  
20 effective amount” refers to that amount of a compound of the present invention which  
lowers IOP when administered to a patient, especially a mammal. The preferred route of  
administration is topical. The compounds of the present invention can be administered as  
solutions, suspensions, or emulsions (dispersions) in an ophthalmically acceptable vehicle.  
As used herein, the term “ophthalmically acceptable vehicle” refers to any substance or  
25 combination of substances which are non-reactive with the compounds and suitable for  
administration to a patient. Solubilizers and stabilizers are deemed to be non-reactive.  
Preferred are aqueous vehicles suitable for topical application to the patient's eyes.

In forming compositions for topical administration, the compounds of the present  
30 invention are generally formulated as between about 0.00003 to about 0.5 percent by  
weight (wt%) solutions in water at a pH between 4.5 to 8.0. The compounds are

preferably formulated as between about 0.0005 to about 0.03 wt% and, most preferably, between about 0.001 and about 0.01 wt%. While the precise regimen is left to the discretion of the clinician, it is recommended that the resulting solution be topically applied by placing one drop in each eye one or two times a day.

5

Other ingredients which may be desirable to use in the ophthalmic preparations of the present invention include preservatives, co-solvents and viscosity building agents.

Antimicrobial Preservatives:

10

Ophthalmic products are typically packaged in multidose form. Preservatives are thus required to prevent microbial contamination during use. Suitable preservatives include: benzalkonium chloride, thimerosal, chlorobutanol, methyl paraben, propyl paraben, phenylethyl alcohol, edetate disodium, sorbic acid, Onamer M, or other agents known to those skilled in the art. Such preservatives are typically employed at a level

15 between about 0.001% and about 1.0% by weight.

Co-Solvents:

20

Prostaglandins, and particularly ester derivatives, typically have limited solubility in water and therefore may require a surfactant or other appropriate co-solvent in the composition. Such co-solvents include: Polysorbate 20, 60 and 80; Pluronic F-68, F-84 and P-103; CREMOPHORE<sup>®</sup> EL (polyoxyl 35 castor oil) cyclodextrin; or other agents known to those skilled in the art. Such co-solvents are typically employed at a level between about 0.01% and about 2% by weight.



Viscosity Agents:

Viscosity greater than that of simple aqueous solutions may be desirable to increase ocular absorption of the active compound, to decrease variability in dispensing the formulations, to decrease physical separation of components of a suspension or emulsion of formulation and/or otherwise to improve the ophthalmic formulation. Such viscosity building agents include, for example, polyvinyl alcohol, polyvinyl pyrrolidone, methyl cellulose, hydroxy propyl methylcellulose, hydroxyethyl cellulose, carboxymethyl cellulose, hydroxy propyl cellulose, chondroitin sulfate and salts thereof, hyaluronic acid and salts thereof, and other agents known to those skilled in the art. Such agents are typically employed at a level between about 0.01% and about 2% by weight

Preferred formulations of *cis*- $\Delta^4$  prostaglandins of the present invention include the following Examples 8-11:

**EXAMPLE 8**

Ingredient	Amount (wt%)
Compound IV or V	0.001
Phosphate Buffered Saline	1.0
Polysorbate 80	0.5
Purified water	q.s. to 100%

**EXAMPLE 9**

	<b>Ingredient</b>	<b>Amount (wt%)</b>
5	Compound IV or V	0.001
	Monobasic sodium phosphate	0.05
	Dibasic sodium phosphate (anhydrous)	0.15
	Sodium chloride	0.75
10	Disodium EDTA (Edetate disodium)	0.05
	Cremophor EL	0.1
	Benzalkonium chloride	0.01
	HCl and/or NaOH	pH 7.3 - 7.4
15	Purified water	q.s. to 100%

**EXAMPLE 10**

	<b>Ingredient</b>	<b>Amount (wt%)</b>
5	Compound IV or V	0.005
	Monobasic sodium phosphate	0.05
	Dibasic sodium phosphate (anhydrous)	0.15
	Sodium chloride	0.75
10	Disodium EDTA (Edetate disodium)	0.05
	CREMOPHOR <sup>®</sup> EL	0.1
	Benzalkonium chloride	0.01
	HCl and/or NaOH	pH 7.3 - 7.4
15	Purified water	q.s. to 100%

**EXAMPLE 11**


	<b>Ingredient</b>	<b>Amount (wt%)</b>
20	Compound IV or V	0.01
	Phosphate Buffered Saline	1.0
	Hydroxypropyl- $\beta$ -cyclodextrin	4.0
25	Purified water	q.s. to 100%

The invention has been described by reference to certain preferred embodiments; however, it should be understood that it may be embodied in other specific forms or variations thereof without departing from its spirit or essential characteristics. The  
30 embodiments described above are therefore considered to be illustrative in all respects and

not restrictive, the scope of the invention being indicated by the appended claims rather than by the foregoing description.

What is claimed is:

1. A method of treating glaucoma or ocular hypertension in a patient, which comprises administering to the patient a pharmaceutically effective amount of a compound of formula **III**:




wherein:

A = CO<sub>2</sub>R, CONR<sup>1</sup>R<sup>2</sup>, CH<sub>2</sub>OR<sup>3</sup>, or CH<sub>2</sub>NR<sup>4</sup>R<sup>5</sup>; where R = H or cationic salt moiety, or CO<sub>2</sub>R = ophthalmically acceptable ester moiety; R<sup>1</sup>, R<sup>2</sup> = same or different = H or alkyl; R<sup>3</sup> = H, acyl, or alkyl; R<sup>4</sup>, R<sup>5</sup> = same or different = H, acyl, or alkyl, with the proviso that if one of R<sup>4</sup>, R<sup>5</sup> = acyl, then the other = H or alkyl;

n = 0 or 2;

L = OR<sup>6</sup> in the α configuration, where R<sup>6</sup> = H, alkyl, or acyl;

B = , where R<sup>7</sup> = H, alkyl, acyl;


---- = single or *trans* double bond;

D, D<sup>1</sup> = different = H and OR<sup>8</sup>, where R<sup>8</sup> = H, alkyl, acyl;

$X = (\text{CH}_2)_m$  or  $(\text{CH}_2)_m\text{O}$ , where  $m = 1-6$ ; and

$Y$  = a phenyl ring optionally substituted with alkyl, halo, trihalomethyl, alkoxy, acyl, acyloxy, amino, alkylamino, acylamino, or hydroxy; or

$X-Y = (\text{CH}_2)_p Y^1$ ; where  $p = 0-6$ ; and



wherein:


$W = \text{CH}_2, \text{O}, \text{S}(\text{O})_q, \text{NR}^9, \text{CH}_2\text{CH}_2, \text{CH}=\text{CH}, \text{CH}_2\text{O}, \text{CH}_2\text{S}(\text{O})_q, \text{CH}=\text{N}$ , or  $\text{CH}_2\text{NR}^9$ ; where  $q = 0-2$ , and  $R^9 = \text{H}$ , alkyl, or acyl;

$Z = \text{H}$ , alkyl, alkoxy, acyl, acyloxy, halo, trihalomethyl, amino, alkylamino, acylamino, or hydroxy; and

---- = single or double bond;

with the proviso that if:

$L = \text{OR}^6$  in the  $\alpha$  configuration, where  $R^6$  is as defined above;



--- = *trans* double bond;

D, D<sup>1</sup> = different = H and OR<sup>8</sup>, where R<sup>8</sup> is as defined above; and

X = CH<sub>2</sub>CH<sub>2</sub> or CH<sub>2</sub>O; then

Y ≠ a phenyl ring, optionally substituted with halo.

2. The method of claim 1, wherein the compound is administered topically.
3. The method of claim 2, wherein the compound is administered as a solution, suspension or emulsion.


4. The method of claim 1, wherein:

A = CO<sub>2</sub>R, where R = H; or CO<sub>2</sub>R = ophthalmically acceptable ester moiety, where R = alkyl;

n = 0;

L = OH in the α configuration;

R<sup>6</sup> = H;




D = OH in the α configuration, and D<sup>1</sup> = H in the β configuration;

X = CH<sub>2</sub>CH<sub>2</sub> or CH<sub>2</sub>O; and

Y = phenyl, optionally substituted with halo or trihalomethyl; or


X-Y =



with the proviso that if:

L = OH in the α configuration;





--- = *trans* double bond;

D = OH in the  $\alpha$  configuration, and D<sup>1</sup> = H in the  $\beta$  configuration; and

X = CH<sub>2</sub>CH<sub>2</sub> or CH<sub>2</sub>O; then


Y  $\neq$  a phenyl ring, optionally substituted with halo.

5. The method of claim 2, wherein the concentration of the compound is between about 0.00003 to about 0.5 weight percent.


6. The method of claim 5, wherein the concentration of the compound is between about 0.0005 to about 0.03 weight percent.

7. The method of claim 6, wherein the concentration of the compound is between about 0.001 to about 0.01 weight percent.


8. The method of claim 5, wherein the compound is:



9. The method of claim 5, wherein the compound is:



10. A compound of formula III:




wherein:

A = CO<sub>2</sub>R, CONR<sup>1</sup>R<sup>2</sup>, CH<sub>2</sub>OR<sup>3</sup>, or CH<sub>2</sub>NR<sup>4</sup>R<sup>5</sup>; where R = H or a cationic salt moiety or CO<sub>2</sub>R forms a pharmaceutically acceptable ester moiety; R<sup>1</sup>, R<sup>2</sup> = same or different = H or alkyl; R<sup>3</sup> = H, acyl, or alkyl; R<sup>4</sup>, R<sup>5</sup> = same or different = H, acyl, or alkyl; with the proviso that if one of R<sup>4</sup>, R<sup>5</sup> = acyl, then the other = H or alkyl;

n = 0 or 2;


L = OR<sup>6</sup> in the α configuration, where R<sup>6</sup> = H, alkyl, or acyl;


  
 B =  $\text{R}^7\text{O}$ , where R<sup>7</sup> = H, alkyl, acyl;

---- = single or *trans* double bond;

D, D<sup>1</sup> = different = H and OR<sup>8</sup>, where R<sup>8</sup> = H, alkyl, acyl;

X-Y = (CH<sub>2</sub>)<sub>p</sub>Y<sup>1</sup>; where p = 0-6; and



wherein:

W = CH<sub>2</sub>, O, S(O)<sub>q</sub>, NR<sup>9</sup>, CH<sub>2</sub>CH<sub>2</sub>, CH=CH, CH<sub>2</sub>O, CH<sub>2</sub>S(O)<sub>q</sub>, CH=N, or CH<sub>2</sub>NR<sup>9</sup>; where q = 0-2, and R<sup>9</sup> = H, alkyl, or acyl;

Z = H, alkyl, alkoxy, acyl, acyloxy, halo, trihalomethyl, amino, alkylamino, acylamino, or hydroxy; and


---- = single or double bond.

11. The compound of claim 10, wherein:

A = CO<sub>2</sub>R, where R = H; or CO<sub>2</sub>R = pharmaceutically acceptable ester moiety,  
where R = or alkyl;


n = 0;

L = OH in the α configuration;




D = OH in the α configuration, and D<sup>1</sup> = H in the β configuration;


X-Y =



12. The compound of claim 11, having the formula:



13. A topical ophthalmic composition for the treatment of glaucoma and ocular hypertension, comprising a compound of formula **III**:




wherein:

A = CO<sub>2</sub>R, CONR<sup>1</sup>R<sup>2</sup>, CH<sub>2</sub>OR<sup>3</sup>, or CH<sub>2</sub>NR<sup>4</sup>R<sup>5</sup>; where R = H or cationic salt moiety, or CO<sub>2</sub>R = ophthalmically acceptable ester moiety; R<sup>1</sup>, R<sup>2</sup> = same or different = H or alkyl; R<sup>3</sup> = H, acyl, or alkyl; R<sup>4</sup>, R<sup>5</sup> = same or different = H, acyl, or alkyl, with the proviso that if one of R<sup>4</sup>, R<sup>5</sup> = acyl, then the other = H or alkyl;

n = 0 or 2;

L = OR<sup>6</sup> in the α configuration, where R<sup>6</sup> = H, alkyl, or acyl;

B = , where R<sup>7</sup> = H, alkyl, acyl;


---- = single or *trans* double bond;

D, D<sup>1</sup> = different = H and OR<sup>8</sup>, where R<sup>8</sup> = H, alkyl, acyl;

X = (CH<sub>2</sub>)<sub>m</sub> or (CH<sub>2</sub>)<sub>m</sub>O, where m = 1-6; and

Y = a phenyl ring optionally substituted with alkyl, halo, trihalomethyl, alkoxy, acyl, acyloxy, amino, alkylamino, acylamino, or hydroxy; or

X-Y = (CH<sub>2</sub>)<sub>p</sub>Y<sup>1</sup>; where p = 0-6; and



wherein:

W = CH<sub>2</sub>, O, S(O)<sub>q</sub>, NR<sup>9</sup>, CH<sub>2</sub>CH<sub>2</sub>, CH=CH, CH<sub>2</sub>O, CH<sub>2</sub>S(O)<sub>q</sub>, CH=N, or CH<sub>2</sub>NR<sup>9</sup>; where q = 0-2, and R<sup>9</sup> = H, alkyl, or acyl;


Z = H, alkyl, alkoxy, acyl, acyloxy, halo, trihalomethyl, amino, alkylamino, acylamino, or hydroxy; and

---- = single or double bond;

and an ophthalmically acceptable vehicle therefor;

with the proviso that if:

L = OR<sup>6</sup> in the α configuration, where R<sup>6</sup> is as defined above;



--- = *trans* double bond;

D, D<sup>1</sup> = different = H and OR<sup>8</sup>, where R<sup>8</sup> is as defined above; and

X = CH<sub>2</sub>CH<sub>2</sub> or CH<sub>2</sub>O; then


Y ≠ a phenyl ring, optionally substituted with halo.

14. The composition of claim 13, wherein:

A = CO<sub>2</sub>R, where R = H; or CO<sub>2</sub>R = ophthalmically acceptable ester moiety, where R = or alkyl;

n = 0;

L = OH in the α configuration;




D = OH in the α configuration, and D<sup>1</sup> = H in the β configuration;

X = CH<sub>2</sub>CH<sub>2</sub> or CH<sub>2</sub>O; and


Y = phenyl, optionally substituted with halo or trihalomethyl; or

X-Y =



with the proviso that if:

L = OH in the  $\alpha$  configuration;




--- = *trans* double bond;

D = OH in the  $\alpha$  configuration, and D' = H in the  $\beta$  configuration; and

X = CH<sub>2</sub>CH<sub>2</sub> or CH<sub>2</sub>O; then

Y  $\neq$  a phenyl ring, optionally substituted with halo.

15. The composition of claim 14, wherein the compound is:



16. The composition of claim 14, wherein the compound is:

